

APC Rabbit anti-Mouse ICOS/CD278 mAb

Catalog No.: A25964

Basic Information

Observed MW

Calculated MW

23KDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC62052-APC

Conjugate

APC. Ex:650nm. Em:660nm.

Background

Predicted to be involved in T cell costimulation; T cell tolerance induction; and cell-cell adhesion. Located in external side of plasma membrane. Is integral component of plasma membrane. Used to study common variable immunodeficiency. Human ortholog(s) of this gene implicated in celiac disease; common variable immunodeficiency; and immunoglobulin alpha deficiency. Orthologous to human ICOS (inducible T cell costimulator).

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Immunogen Information

Gene ID

54167

Swiss Prot

Q9WVS0

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 21-142 of mouse ICOS/CD278(NP_059508.2).

Synonyms

H4; CCLP; AILIM; CRP-1; Ly115

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification



Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data



Flow cytometry: 1X10⁶ CHO cells (negative control, left) and CHO (Transfection, right) cells were surface-stained with APC Rabbit anti-Mouse ICOS/CD278 mAb (A25964, 5 µl/Test, orange line) or APC Rabbit IgG isotype control (A24173, 5 µl/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10⁶ CHO (Transfection) cells were surface-stained with APC Rabbit IgG isotype control (A24173, 5 µl/Test, left) or APC Rabbit anti-Mouse ICOS/CD278 mAb (A25964, 5 µl/Test, right).