

# ABflo® 700 Rabbit anti-Human/Monkey CD20 mAb

Catalog No.: A25966

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

14kDa/33kDa

### Category

Primary antibody

### Applications

FC

### Cross-Reactivity

Human, Cynomolgus

### CloneNo number

ARC51683-04

### Conjugate

ABflo® 700. Ex:690nm. Em:713nm.

## Recommended Dilutions

**FC** 5 µl per 10<sup>6</sup> cells in  
100 µl volume

## Background

This gene encodes a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. This gene encodes a B-lymphocyte surface molecule which plays a role in the development and differentiation of B-cells into plasma cells. This family member is localized to 11q12, among a cluster of family members. Alternative splicing of this gene results in two transcript variants which encode the same protein.

## Immunogen Information

### Gene ID

931

### Swiss Prot

P11836

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

B1; S7; Bp35; CD20; FMC7; CVID5; LEU-16

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Validation Data



Flow cytometry:  $1 \times 10^6$  Jurkat cells (negative control, left) and Daudi cells (right) were surface-stained with ABflo® 700 Rabbit anti-Human/Monkey CD20 mAb (A25966, 5  $\mu$ l/Test, orange line) or ABflo® 700 Rabbit IgG isotype control (A25976, 5  $\mu$ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:  $1 \times 10^6$  Daudi cells were surface-stained with ABflo® 700 Rabbit IgG isotype control (A25976, 5  $\mu$ l/Test, left) or ABflo® 700 Rabbit anti-Human/Monkey CD20 mAb (A25966, 5  $\mu$ l/Test, right).



Flow cytometry:  $1 \times 10^6$  Cynomolgus Monkey PBMC were surface-stained with PE Rabbit anti-Human/Monkey CD19 mAb (A26609, 5  $\mu$ l/Test) and ABflo® 700 Rabbit IgG isotype control (A25976, 5  $\mu$ l/Test, left) or ABflo® 700 Rabbit anti-Human/Monkey CD20 mAb (A25966, 5  $\mu$ l/Test, right). Cells in the lymphocyte gate were used for analysis.