Magnetic Beads-conjugated Pan Acetyl-Lysine Rabbit mAb

Catalog No.: A26013 Recombinant

Basic Information

Observed MW

Calculated MW

Category Primary antibody

Applications

Cross-Reactivity Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number ARC67569-MB

Conjugate

Magnetic Beads

Recommended Dilutions

IP

30ul antibody (bead slurry) for 200µg-400µg extracts of whole cells

Background

Acetylation of lysine, like phosphorylation of serine, threonine or tyrosine, is an important reversible modification controlling protein activity. The conserved amino-terminal domains of the four core histones (H2A, H2B, H3, and H4) contain lysines that are acetylated by histone acetyltransferases (HATs) and deacetylated by histone deacetylases (HDACs) . Signaling resulting in acetylation/deacetylation of histones, transcription factors, and other proteins affects a diverse array of cellular processes including chromatin structure and gene activity, cell growth, differentiation, and apoptosis . Recent proteomic surveys suggest that acetylation of lysine residues may be a widespread and important form of post-translational protein modification that affects thousands of proteins involved in control of cell cycle and metabolism, longevity, actin polymerization, and nuclear transport . The regulation of protein acetylation status is impaired in cancer and polyglutamine diseases, and HDACs have become promising targets for anti-cancer drugs currently in development .

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Contact

6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn
Θ	www.abclonal.com.cn

Product Information

Source Rabbit **Isotype** IgG **Purification** Affinity purification

Storage

Store at 4°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,pH7.3.

