

Magnetic Beads-conjugated Pan Succinyl-Lysine Rabbit mAb

www.abclonal.comCatalog No.: A26017 **Recombinant**

Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

IP

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC55831-MB

Conjugate

Magnetic Beads

Background

The lysine residue in proteins can undergo to many types of PTMs, such as methylation, acetylation, biotinylation, ubiquitination, ubiquitin-like modifications, propionylation and butyrylation, then leading to the remarkable complexity of PTM networks. Succinylation is a posttranslational modification (PTM) where a succinyl group is added to a Lys (K) residue of a protein molecule. It has been shown that the lysine succinylation responses to different physiological conditions and is evolutionary conserved. It has been identified 2,565 succinylation sites from 779 proteins and revealed that lysine succinylation have potential impacts on enzymes involved in mitochondrial metabolism including the amino acid degradation, tricarboxylic acid (TCA) cycle and fatty acid metabolism. The lysine succinylation also occurs in histones, suggesting that it may play an important role in regulating chromatin structures and functions as well.

Recommended Dilutions

IP 30ul antibody (bead slurry) for 200µg-400µg extracts of whole cells

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at 4°C. Avoid freeze / thaw cycles.
Buffer: PBS with 0.05% proclin300, pH7.3.