

APC Rabbit anti-Mouse CD138/Syndecan-1 mAb

Catalog No.: A26049

Basic Information

Observed MW

Calculated MW

33kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC66008-APC

Conjugate

APC. Ex:650nm. Em:660nm.

Background

Predicted to enable identical protein binding activity and protein C-terminus binding activity. Involved in myoblast development. Acts upstream of or within canonical Wnt signaling pathway. Located in external side of plasma membrane. Is expressed in several structures, including alimentary system; egg cylinder; sensory organ; skin; and urinary system. Orthologous to human SDC1 (syndecan 1).

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 20969

Swiss Prot

P18828

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 23-255 of mouse CD138/Syndecan-1 (NP_035649.1).

Synonyms

Sstn; Synd; CD138; Synd1; syn-1; CD138/Syndecan-1

Contact

2		400-999-6126
\bowtie	Τ	cn.market@abclonal.com.cn
•	Т	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data





Flow cytometry: 1X10^6 C57BL/6 mouse bone marrow cells were surface-stained with APC Rabbit IgG isotype control (A24173,5 µl/Test,left) or APC Rabbit anti-Mouse CD138/Syndecan-1 mAb (A26049,5 µl/Test,right).

Flow cytometry: 1X10^6 C57BL/6 mouse bone marrow cells were surface-stained with ABflo® 488 Rabbit anti-Mouse CD19 mAb (A24100,5 µl/Test) and APC Rabbit anti-Mouse CD138/Syndecan-1 mAb (A26049,5 µl/Test,right) or APC Rabbit IgG isotype control (A24173,5 µl/Test,left).