

APC Rabbit anti-Mouse CD93 mAb

Catalog No.: A26058

Basic Information

Observed MW

Refer to figures

Calculated MW

69kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC67562

Conjugate

APC. Ex:650nm. Em:660nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

Predicted to enable complement component C1q complex binding activity. Predicted to be involved in cell-cell adhesion. Predicted to act upstream of or within cell adhesion. Located in cell surface; cytoplasmic vesicle; and plasma membrane. Is expressed in blood vessel. Orthologous to human CD93 (CD93 molecule).

Immunogen Information

Gene ID

17064

Swiss Prot

O89103

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Ly68; AA4.1; C1qr1; C1qrp; 6030404G09Rik

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.


Validation Data



Flow cytometry: 1×10^6 293T cells (negative control, left) and 293T (Transfection, right) cells were surface-stained with APC Rabbit anti-Mouse CD93 mAb (A26058,5 $\mu\text{l}/\text{Test}$, orange line) or APC Rabbit IgG isotype control (A24173,5 $\mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 293T (Transfection) cells were surface-stained with APC Rabbit IgG isotype control (A24173,5 $\mu\text{l}/\text{Test}$, left) or APC Rabbit anti-Mouse CD93 mAb (A26058,5 $\mu\text{l}/\text{Test}$, right).



Flow cytometry: 1×10^6 C57BL/6 mouse bone marrow cells were surface-stained with APC Rabbit anti-Mouse CD93 mAb (A26058,5 $\mu\text{l}/\text{Test}$, orange line) or APC Rabbit IgG isotype control (A24173,5 $\mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line).