Leader in Biomolecular Solutions for Life Science

PE Rabbit anti-Human CD170/Siglec-5 mAb

Catalog No.: A26119



Basic Information

Observed MW

Calculated MW 61kDa

Category Primary antibody

Applications FC

Cross-Reactivity Human

CloneNo number ARC60966-PE

Conjugate

PE. Ex:565nm. Em:574nm.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Background

This gene encodes a member of the sialic acid-binding immunoglobulin-like lectin (Siglec) family. These cell surface lectins are characterized by structural motifs in the immunoglobulin (Ig)-like domains and sialic acid recognition sites in the first Ig V set domain. The encoded protein is a member of the CD33-related subset of Siglecs and inhibits the activation of several cell types including monocytes, macrophages and neutrophils. Binding of group B Streptococcus (GBS) to the encoded protein plays a role in GBS immune evasion.

Immunogen Information

Gene ID 8778

Swiss Prot 015389

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 17-434 of human CD170/Siglec-5 (NP_003821.1).

Synonyms

CD170; OBBP2; CD33L2; OB-BP2; SIGLEC-5

Contact

400-999-6126 <u>cn.market@abclonal.com.cn</u> <u>www.abclonal.com.cn</u>


Product Information


Source Rabbit **Isotype** IgG **Purification** Affinity purification


Storage

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.

Validation Data







Flow cytometry: 1X10^6 K-562 cells (negative control,left) and U-937 cells (right) were surface-stained with PE Rabbit anti-Human CD170/Siglec-5 mAb (A26119,5 µl/Test,orange line) or PE Rabbit IgG isotype control (A24172,5 µl/Test,blue line). Nonfluorescently stained cells were used as blank control (red line). Flow cytometry: 1X10^6 U-937 cells were surface-stained with PE Rabbit IgG isotype control (A24172,5 μ //Test,left) or PE Rabbit anti-Human CD170/Siglec-5 mAb (A26119,5 μ //Test,right).

Flow cytometry: 1X10^6 Human PBMC were surface-stained with PE Rabbit anti-Human CD170/Siglec-5 mAb (A26119,5 µl/Test,orange line) or PE Rabbit IgG isotype control (A24172,5 µl/Test,blue line). Nonfluorescently stained cells were used as blank control (red line).