Leader in Biomolecular Solutions for Life Science

# APC Rabbit anti-Mouse CD32 mAb

Catalog No.: A26143



## **Basic Information**

**Observed MW** Refer to figures

Calculated MW 31kDa/33kDa/36kDa

Category Primary antibody

Applications FC

Cross-Reactivity Mouse

CloneNo number ARC68229-APC

Conjugate

APC. Ex:650nm. Em:660nm.

## **Recommended Dilutions**

5 µl per 10^6 cells in

100 µl volume

FC

## **Immunogen Information**

**Gene ID** 14130 Swiss Prot P08101

#### Immunogen

Background

Recombinant fusion protein containing a sequence corresponding to amino acids 30-207 of mouse CD32 (NP\_034317.1).

CD32 (Fcgr2) is a 40 kD transmembrane glycoprotein, member of the immunoglobulin superfamily. The extracellular region of CD32 consists of two Ig C-type domains that binds

expressed by Macrophages, neutrophils, mast cells and B cells.

the Fc region from monomeric IgG with low affinity, but binds immune complexes efficiently. CD32 can mediate phagocytosis of immune complexes and modulate cell activation. CD32 is

Synonyms

CD32; Fcgr2; Fcr-2; Fcr-3; Ly-17; LyM-1; Lym-1; fcRII; FcgRII; Fcgr2a; Ly-m20; Fc[g]RII; F630109E10Rik

## Contact

6	400-999-6126
$\times$	cn.market@abclonal.com.cn
€	www.abclonal.com.cn


## **Product Information**


**Source** Rabbit **lsotype** IgG Purification Affinity purification


### Storage

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.

## Validation Data







Flow cytometry: 1X10^6 bEnd.3 cells (negative control,left) and C57BL/6 mouse splenocytes (right) were surface-stained with APC Rabbit anti-Mouse CD32 mAb (A26143,5 µl/Test,orange line) or APC Rabbit IgG isotype control (A24173,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line). Flow cytometry: 1X10^6 C57BL/6 mouse splenocytes were surface-stained with APC Rabbit IgG isotype control (A24173,5  $\mu$ I/Test,left) or APC Rabbit anti-Mouse CD32 mAb (A26143,5  $\mu$ I/Test,right).

Flow cytometry:1X10^6 C57BL/6 mouse splenocytes were surface-stained with PE Rabbit anti-Mouse CD19 mAb (A25265,5 µl/Test) and APC Rabbit anti-Mouse CD32 mAb (A26143,5 µl/Test,right) or APC Rabbit IgG isotype control (A24173,5 µl/Test,left).