# ABflo® 450 Rabbit anti-Mouse CD38 mAb

ABclonal www.abclonal.com

Catalog No.: A26194

## **Basic Information**

#### **Observed MW**

Refer to figures

### **Calculated MW**

34kDa

### Category

Primary antibody

### **Applications**

FC

### **Cross-Reactivity**

Mouse

#### CloneNo number

ARC59835-ABflo450

### Conjugate

ABflo® 450. Ex:406nm. Em:445nm.

## **Background**

This gene encodes a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Knockout mice deficient for this gene display altered humoral immune responses. In addition, knockout mice exhibit higher locomotor activity and defects in nurturing and social behaviors.

## **Recommended Dilutions**

FC

5 μl per 10^6 cells in 100 μl volume

## **Immunogen Information**

**Gene ID** 12494

**Swiss Prot** 

P56528

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 45-304 of mouse CD38 (NP\_031672.2).

## **Synonyms**

I-19; ADPRC 1; Cd38-rs1

## **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
<u>~</u>	www.abclonal.com.cn

## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

## **Validation Data**









Flow cytometry: 1X10^6 C2C12 cells (negative control,left) and C57BL/6 mouse splenocytes (right) were surface-stained with ABflo® 450 Rabbit anti-Mouse CD38 mAb (A26194,5 µl/Test,orange line) or ABflo® 450 Rabbit IgG isotype control (5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1X10^6$  C57BL/6 mouse splenocytes were surface-stained with ABflo® 450 Rabbit IgG isotype control (5  $\mu$ I/Test,left) or ABflo® 450 Rabbit anti-Mouse CD38 mAb (A26194,5  $\mu$ I/Test,right).