ABclonal www.abclonal.com

APC/Cyanine7 Mouse anti-Human CD3 mAb

Catalog No.: A26219

Basic Information

Observed MW

Calculated MW

23kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

Conjugate

APC-Cy7. Ex:651nm. Em:779nm.

Background

The protein encoded by this gene is the CD3-epsilon polypeptide, which together with CD3-gamma, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. Defects in this gene cause immunodeficiency. This gene has also been linked to a susceptibility to type I diabetes in women.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 916 Swiss Prot P07766

Immunogen

Recombinant protein of human CD3.

Synonyms

T3E; TCRE; IMD18; CD3epsilon

Contact

2		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


Product Information


SourceIsotypePurificationMouseIqGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.





1X10^6 Human PBMC were surface-stained with APC/Cyanine7 Mouse anti-Human CD3 mAb (A26219,5 μ l/Test,orange line) or APC/Cyanine7 Rabbit IgG isotype control (5 μ l/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

1X10^6 Human PBMC were surface-stained with APC/Cyanine7 Rabbit IgG isotype control (5 μ I/Test,left) or APC/Cyanine7 Mouse anti-Human CD3 mAb (A26219,5 μ I/Test,right).