

ABflo® 700 Rabbit anti-Human ITGA1/CD49a mAb

Catalog No.: A26235

Basic Information

Observed MW

Refer to figures

Calculated MW

131kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC58076

Conjugate

ABflo® 700. Ex:690nm. Em:713nm.

Background

This gene encodes the alpha 1 subunit of integrin receptors. This protein heterodimerizes with the beta 1 subunit to form a cell-surface receptor for collagen and laminin. The heterodimeric receptor is involved in cell-cell adhesion and may play a role in inflammation and fibrosis. The alpha 1 subunit contains an inserted (I) von Willebrand factor type I domain which is thought to be involved in collagen binding.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 3672 **Swiss Prot**

P56199

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

VLA1; CD49a

Contact

a	400-999-6126
\bowtie	cn.market@abclonal.com.cn
<u></u>	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1X10^6 K-562 cells (negative control,left) and SH-SY5Y cells (right) were surface-stained with ABflo® 700 Rabbit anti-Human ITGA1/CD49a mAb (A26235,5 µl/Test,orange line) or ABflo® 700 Rabbit IgG isotype control (A25976,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).