

# ABflo® 700 Rabbit anti-Human CD271/NGFR mAb

Catalog No.: A26369

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

35kDa/45kDa

### Category

Primary antibody

### Applications

FC

### Cross-Reactivity

Human

### CloneNo number

ARC57800-ABflo700

### Conjugate

ABflo® 700. Ex:690nm. Em:713nm.

## Recommended Dilutions

**FC** 5 µl per 10<sup>6</sup> cells in  
100 µl volume

## Background

Nerve growth factor receptor contains an extracellular domain containing four 40-amino acid repeats with 6 cysteine residues at conserved positions followed by a serine/threonine-rich region, a single transmembrane domain, and a 155-amino acid cytoplasmic domain. The cysteine-rich region contains the nerve growth factor binding domain.

## Immunogen Information

### Gene ID

4804

### Swiss Prot

P08138

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 29-250 of human CD271/NGFR (NP\_002498.1).

### Synonyms

CD271; p75NTR; TNFRSF16; p75(NTR); Gp80-LNGFR

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

## Validation Data



Flow cytometry:  $1 \times 10^6$  K-562 cells (negative control, left) and SK-N-MC cells (right) were surface-stained with ABflo® 700 Rabbit anti-Human NGFR/CD271 mAb (A26369, 5  $\mu$ l/Test, orange line) or ABflo® 700 Rabbit IgG isotype control (A25976, 5  $\mu$ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1 \times 10^6$  SK-N-MC cells were surface-stained with ABflo® 700 Rabbit IgG isotype control (A25976, 5  $\mu$ l/Test, left) or ABflo® 700 Rabbit anti-Human NGFR/CD271 mAb (A26369, 5  $\mu$ l/Test, right).