

ABflo® 700 Rabbit anti-Human CD271/NGFR mAb

Catalog No.: A26369

Basic Information

Observed MW

Refer to figures

Calculated MW

35kDa/45kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC57800-ABflo700

Conjugate

ABflo® 700. Ex:690nm. Em:713nm.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
 100 µl volume

Background

Nerve growth factor receptor contains an extracellular domain containing four 40-amino acid repeats with 6 cysteine residues at conserved positions followed by a serine/threonine-rich region, a single transmembrane domain, and a 155-amino acid cytoplasmic domain. The cysteine-rich region contains the nerve growth factor binding domain.

Immunogen Information

Gene ID

4804

Swiss Prot

P08138

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 29-250 of human CD271/NGFR (NP_002498.1).

Synonyms

CD271; p75NTR; TNFRSF16; p75(NTR); Gp80-LNGFR

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 K-562 cells (negative control, left) and SK-N-MC cells (right) were surface-stained with ABflo® 700 Rabbit anti-Human NGFR/CD271 mAb (A26369, 5 μ l/Test, orange line) or ABflo® 700 Rabbit IgG isotype control (A25976, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 SK-N-MC cells were surface-stained with ABflo® 700 Rabbit IgG isotype control (A25976, 5 μ l/Test, left) or ABflo® 700 Rabbit anti-Human NGFR/CD271 mAb (A26369, 5 μ l/Test, right).