

PE Rabbit anti-Mouse CD68 mAb

Catalog No.: A26390

Basic Information

Observed MW

Calculated MW

35kDa

Category

Primary antibody

Applications

FC (intra)

Cross-Reactivity

Mouse

CloneNo number

ARC60987-PE

Conjugate

PE. Ex:565nm. Em:574nm.

Recommended Dilutions

FC (intra) 5 μ l per 10^6 cells in
100 μ l volume

Background

Involved in several processes, including cellular response to lipopolysaccharide; cellular response to oxidised low-density lipoprotein particle stimulus; and negative regulation of dendritic cell antigen processing and presentation. Acts upstream of or within aging and cellular response to organic substance. Located in lysosome and plasma membrane. Is expressed in several structures, including central nervous system; embryo mesenchyme; heart blood vessel; liver; and spleen. Orthologous to human CD68 (CD68 molecule).

Immunogen Information

Gene ID

12514

Swiss Prot

P31996

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 21-291 of mouse CD68 (NP_001277987.1).

Synonyms

Lamp4; gp110; Scard1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 NIH/3T3 cells (Low Expression, left) and RAW 264.7 cells (right) were intracellularly-stained with PE Rabbit anti-Mouse CD68 mAb (A26390, 5 μ l/Test, orange line) or PE Rabbit IgG isotype control (A24172, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 RAW 264.7 cells were intracellularly-stained with PE Rabbit IgG isotype control (A24172, 5 μ l/Test, left) or PE Rabbit anti-Mouse CD68 mAb (A26390, 5 μ l/Test, right).