Leader in Biomolecular Solutions for Life Science

PE Rabbit anti-Mouse CD68 mAb

Catalog No.: A26390



Basic Information

Observed MW

Calculated MW 35kDa

Category Primary antibody

Applications FC (intra)

Cross-Reactivity Mouse

CloneNo number ARC60987

Conjugate PE. Ex:565nm. Em:574nm.

Recommended Dilutions

FC (intra)

5 μl per 10^6 cells in 100 μl volume

Background

Involved in several processes, including cellular response to lipopolysaccharide; cellular response to oxidised low-density lipoprotein particle stimulus; and negative regulation of dendritic cell antigen processing and presentation. Acts upstream of or within aging and cellular response to organic substance. Located in lysosome and plasma membrane. Is expressed in several structures, including central nervous system; embryo mesenchyme; heart blood vessel; liver; and spleen. Orthologous to human CD68 (CD68 molecule).

Immunogen Information

Gene ID 12514 Swiss Prot P31996

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Lamp4; gp110; Scard1

Contact

6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn


Product Information


Source Rabbit **lsotype** IgG **Purification** Affinity purification

Storage

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.

Validation Data





Flow cytometry: 1X10^6 NIH/3T3 cells (Low Expression,left) and RAW 264.7 cells (right) were intracellularly-stained with PE Rabbit anti-Mouse CD68 mAb (A26390,5 µl/Test,orange line) or PE Rabbit IgG isotype control (A24172,5 µl/Test,blue line). Nonfluorescently stained cells were used as blank control (red line). Flow cytometry: 1X10^6 RAW 264.7 cells were intracellularly-stained with PE Rabbit IgG isotype control (A24172,5 µl/Test,left) or PE Rabbit anti-Mouse CD68 mAb (A26390,5 µl/Test,right).