

APC Rabbit anti-Human IgG (Fc) mAb

Catalog No.: A26395

Basic Information

Observed MW

Refer to figures

Calculated MW

36kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC53383

Conjugate

APC. Ex:650nm. Em:660nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

Predicted to enable antigen binding activity and immunoglobulin receptor binding activity. Predicted to be involved in several processes, including activation of immune response; defense response to other organism; and phagocytosis. Predicted to act upstream of or within several processes, including immunoglobulin mediated immune response; positive regulation of hypersensitivity; and positive regulation of phagocytosis. Located in extracellular space.

Immunogen Information

Gene ID

3500

Swiss Prot

P01857

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

IGHG1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 Human PBMC were intracellularly-stained with FITC Mouse anti-Human CD19 mAb (A22815, 5 μ l/Test) and APC Rabbit IgG isotype control (A24173, 5 μ l/Test, left) or APC Rabbit anti-Human IgG (Fc) mAb (A26395, 5 μ l/Test, right).