Leader in Biomolecular Solutions for Life Science

APC Rabbit anti-Human CD30 mAb

Catalog No.: A26397



Basic Information

Observed MW

Calculated MW 14kDa/50kDa/63kDa

Category Primary antibody

Applications FC

Cross-Reactivity Human

CloneNo number ARC52955-APC

Conjugate APC. Ex:650nm. Em:660nm.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Immunogen Information

Gene ID 943 Swiss Prot P28908

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 19-379 of human CD30 (NP_001234.3).

Synonyms

CD30; Ki-1; D1S166E

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
€	www.abclonal.com.cn


Product Information


Source Rabbit **Isotype** IgG **Purification** Affinity purification

Storage

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.

Validation Data





Flow cytometry: 1X10^6 A549 cells (negative control,left) and K-562 cells (right) were surface-stained with APC Rabbit anti-Human CD30 mAb (A26397,5 μ l/Test,orange line) or APC Rabbit IgG isotype control (A24173,5 μ l/Test,blue line). Nonfluorescently stained cells were used as blank control (red line). Flow cytometry: 1X10⁶ K-562 cells were surface-stained with APC Rabbit IgG isotype control (A24173,5 µl/Test,left) or APC Rabbit anti-Human CD30 mAb (A26397,5 µl/Test,right).