

ABflo® 488 Rabbit anti-Mouse CD94 mAb

Catalog No.: A26478

Basic Information

Observed MW

Refer to figures

Calculated MW

21kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC67768-ABflo488

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Background

CD94 is a 43/39 kD C-type lectin, also known as Kp43. It is present on all NK cells, NKT cells, and a subset of CD8-positive T lymphocytes in most mouse strains. CD94 is a type-II transmembrane protein with an extracellular lectin-like domain and a short cytoplasmic tail. CD94 is expressed as a disulphide-linked heterodimer with a NKG2 subunit believed to mediate signal transduction. When associated with NKG2A, the complex triggers inhibition; when associated with NKG2C, the complex triggers stimulation. The receptor complex of CD94 and NKG2 receptors bind to the ligand, Qa-1, and are thought to play a role in maintaining self-tolerance in developing NK cells.

Immunogen Information

Gene ID

16643

Swiss Prot

O54707

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 59-179 of mouse CD94 (NP_034784.1).

Synonyms

CD94

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 C57BL/6 mouse splenocytes were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 μ l/Test,left) or ABflo® 488 Rabbit anti-Mouse CD94 mAb (A26478,5 μ l/Test,right).

Flow cytometry: 1×10^6 C57BL/6 mouse splenocytes were surface-stained with ABflo® 647 Rabbit anti-Mouse CD161c/NK1.1 mAb (5 μ l/Test) and ABflo® 488 Rabbit IgG isotype control (A22069,5 μ l/Test,left) or ABflo® 488 Rabbit anti-Mouse CD94 mAb (A26478,5 μ l/Test,right).