

# ABflo® 610 Rabbit anti-Human/Monkey IgD mAb

Catalog No.: A26481

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

47kDa

### Category

Primary antibody

### Applications

FC

### Cross-Reactivity

Human, Rhesus

### CloneNo number

ARC60051

### Conjugate

ABflo® 610. Ex:421nm. Em:612nm.

## Recommended Dilutions

**FC** 5 µl per 10<sup>6</sup> cells in  
100 µl volume

## Background

IgD is the major antigen receptor isotype on the surface of most peripheral B-cells, where it is coexpressed with IgM. The membrane-bound IgD (mIgD) induces the phosphorylation of CD79A and CD79B by the Src family of protein tyrosine kinases.

## Immunogen Information

### Gene ID

3495

### Swiss Prot

P01880

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

IGHD

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.


## Validation Data



Flow cytometry:  $1 \times 10^6$  Jurkat cells (negative control, left) and Human PBMC (right) were surface-stained with ABflo® 610 Rabbit anti-Human IgD mAb (A26481,5  $\mu\text{l}/\text{Test}$ , orange line) or ABflo® 610 Rabbit IgG isotype control (A25826,5  $\mu\text{l}/\text{Test}$ , blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:  $1 \times 10^6$  Human PBMC were surface-stained with ABflo® 610 Rabbit IgG isotype control (A25826,5  $\mu\text{l}/\text{Test}$ , left) or ABflo® 610 Rabbit anti-Human IgD mAb (A26481,5  $\mu\text{l}/\text{Test}$ , right).



Flow cytometry:  $1 \times 10^6$  Rhesus PBMC were surface-stained with ABflo® 488 Mouse anti-Human CD11b mAb (A28074,5  $\mu\text{l}/\text{Test}$ ), ABflo® 450 Rabbit anti-Human/Monkey CD19 mAb (A27286,5  $\mu\text{l}/\text{Test}$ ) and ABflo® 610 Rabbit IgG isotype control (A25826,5  $\mu\text{l}/\text{Test}$ , left) or ABflo® 610 Rabbit anti-Human/Monkey IgD mAb (A26481,5  $\mu\text{l}/\text{Test}$ , right). Cells in the CD11b- gate were used for analysis.