

APC Rabbit anti-Human CD261/TRAIL-R1 mAb

Catalog No.: A26511

Basic Information

Observed MW

Calculated MW

50kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC63581-APC

Conjugate

APC. Ex:650nm. Em:660nm.

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Immunogen Information

Gene ID

8797

Swiss Prot

O00220

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 124-232 of human CD261/TRAIL-R1 (NP_003835.3).

Synonyms

TNFRSF10A; APO2; CD261; DR4; TRAILR-1; TRAILR1; TNF receptor superfamily member 10a

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 BeWo cells (negative control, left) and HeLa cells (right) were surface-stained with APC Rabbit anti-Human CD261/TRAIL-R1 mAb (A26511, 5 μ l/Test, orange line) or APC Rabbit IgG isotype control (A24173, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 HeLa cells were surface-stained with APC Rabbit IgG isotype control (A24173, 5 μ l/Test, left) or APC Rabbit anti-Human CD261/TRAIL-R1 mAb (A26511, 5 μ l/Test, right).