

ABflo® 700 Rabbit anti-Mouse CD5 mAb

Catalog No.: A26557

Basic Information

Observed MW

Calculated MW

54kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC61614-ABflo700

Conjugate

ABflo® 700. Ex:690nm. Em:713nm.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Background

CD5 is a type I transmembrane protein belonging to the scavenger receptor cysteine-enriched (SRCR) family, characterized by the presence of at least one SRCR domain containing 90-110 amino acids. CD5 is expressed in all mature T cells, B-1A subsets of mature B cells, and certain leukemia B cells. It is elevated in regulatory T and B cells (Tregs/Bregs). CD5 expression was also elevated in incompetent T and B cells. Elevated levels of CD5 have also been found in many autoimmune diseases. CD5 binds to T cell receptors (TCR) and negatively regulates T cell activation and differentiation. The expression of CD5 in tumor-infiltrating T lymphocytes was negatively correlated with their antitumor activity. Recently, it has been reported that CD5 binds directly to IL6 and mediates downstream signal transduction. CD5+ B cells promote tumor growth in animal models. Agents targeting CD5 are being actively explored as therapeutic interventions for cancer and other conditions.

Immunogen Information

Gene ID

12507

Swiss Prot

P13379

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 25-371 of mouse CD5 (NP_031676.3).

Synonyms

Ly-1; Ly-A; Ly-12; Lyt-1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification



Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1X10⁶ A20 cells (negative control, left) and CTLL-2 cells (right) were surface-stained with ABflo® 700 Rabbit anti-Mouse CD5 mAb (A26557,5 µl/Test, orange line) or ABflo® 700 Rabbit IgG isotype control (A25976,5 µl/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10⁶ CTLL-2 cells were surface-stained with ABflo® 700 Rabbit IgG isotype control (A25976,5 µl/Test, left) or ABflo® 700 Rabbit anti-Mouse CD5 mAb (A26557,5 µl/Test, right).