

APC Rabbit anti-Mouse CD304/Neuropilin-1 mAb

Catalog No.: A26587

Basic Information

Observed MW

Refer to figures

Calculated MW

103kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC69115

Conjugate

APC. Ex:650nm. Em:660nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

Enables protein kinase binding activity; transmembrane signaling receptor activity; and vascular endothelial growth factor binding activity. Involved in several processes, including nervous system development; regulation of signal transduction; and vasculature development. Acts upstream of or within several processes, including morphogenesis of a branching epithelium; nervous system development; and toxin transport. Located in several cellular components, including endosome; focal adhesion; and neurofilament. Is integral component of postsynaptic membrane. Is expressed in several structures, including cardiovascular system; jaw; lung; nervous system; and sensory organ. Used to study retinal vein occlusion. Human ortholog(s) of this gene implicated in lung non-small cell carcinoma. Orthologous to human NRP1 (neuropilin 1).

Immunogen Information

Gene ID

18186

Swiss Prot

P97333

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Nrp; NP-1; Npn1; NPN-1; C530029I03

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification



Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 CHO cells (negative control, left) and CHO (Transfection, right) cells were surface-stained with APC Rabbit anti-Mouse CD304/Neuropilin-1 mAb ($2.6587, 5 \mu\text{l}/\text{Test}$, orange line) or APC Rabbit IgG isotype control ($2.4173, 5 \mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 C57BL/6 mouse splenocytes were surface-stained with APC Rabbit IgG isotype control ($2.4173, 5 \mu\text{l}/\text{Test}$, left) or APC Rabbit anti-Mouse CD304/Neuropilin-1 mAb ($2.6587, 5 \mu\text{l}/\text{Test}$, right).