

PE Rabbit anti-Mouse CD49a/ITGA1 mAb

Catalog No.: A26653

Basic Information

Observed MW

Refer to figures

Calculated MW

130kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC68868

Conjugate

PE. Ex:565nm. Em:574nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

CD49a is a 1179 aa, type I transmembrane glycoprotein also known as α 1 integrin, VLA-1 α chain, or integrin α 1. It associates antibody v042010 with CD29 (β 1 integrin) to form the VLA-1 complex, a collagen IV and alminin-1 receptor that is expressed on activated T cells, smooth muscle cells, endothelial cells, neuronal cells, fibroblasts, and mesenchymal cells. CD49a is an adhesion molecule and is involved in the regulation of leukocyte migration, T cell proliferation, and cytokine production.

Immunogen Information

Gene ID

109700

Swiss Prot

Q3V3R4

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Vla1; CD49A; E130012M19Rik

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification



Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1X10⁶ NIH/3T3 cells (negative control, left) and Neuro-2a cells (right) were surface-stained with PE Rabbit anti-Mouse Integrin alpha 1/CD49a mAb (A26653, 5 µl/Test, orange line) or PE Rabbit IgG isotype control (A24172, 5 µl/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10⁶ Neuro-2a cells were surface-stained with PE Rabbit IgG isotype control (A24172, 5 µl/Test, left) or PE Rabbit anti-Mouse Integrin alpha 1/CD49a mAb (A26653, 5 µl/Test, right).