

ABflo® 610 Rabbit anti-Human CD94 mAb

Catalog No.: A26683

Basic Information

Observed MW

Calculated MW

21kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC60983-ABflo610

Conjugate

ABflo® 610. Ex:421nm. Em:612nm.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Background

Natural killer (NK) cells are a distinct lineage of lymphocytes that mediate cytotoxic activity and secrete cytokines upon immune stimulation. Several genes of the C-type lectin superfamily, including members of the NKG2 family, are expressed by NK cells and may be involved in the regulation of NK cell function. KLRD1 (CD94) is an antigen preferentially expressed on NK cells and is classified as a type II membrane protein because it has an external C terminus. Several transcript variants encoding different isoforms have been found for this gene.

Immunogen Information

Gene ID

3824

Swiss Prot

Q13241

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 32-179 of human CD94. (NP_002253.2).

Synonyms

CD94

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.


Validation Data




Flow cytometry: 1X10⁶ HAP1 cells (negative control, left) and NK-92MI cells (right) were surface-stained with ABflo® 610 Rabbit anti-Human CD94 mAb (A26683,5 µl/Test, orange line) or ABflo® 610 Rabbit IgG isotype control (A25826,5 µl/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10⁶ NK-92MI cells were surface-stained with ABflo® 610 Rabbit IgG isotype control (A25826,5 µl/Test, left) or ABflo® 610 Rabbit anti-Human CD94 mAb (A26683,5 µl/Test, right).



Flow cytometry: 1X10⁶ Human PBMC were surface-stained with ABflo® 610 Rabbit anti-Human CD94 mAb (A26683,5 µl/Test, orange line) or ABflo® 610 Rabbit IgG isotype control (A25826,5 µl/Test, blue line). Non-fluorescently stained cells were used as blank control (red line). Cells in the Lymphocytes gate were used for analysis.



Flow cytometry: 1X10⁶ Human PBMC were surface-stained with ABflo® 610 Rabbit IgG isotype control (A25826,5 µl/Test, left) or ABflo® 610 Rabbit anti-Human CD94 mAb (A26683,5 µl/Test, right). Cells in the Lymphocytes gate were used for analysis.