DARPP32 Rabbit mAb

Catalog No.: A26749 Recombinant



Basic Information

Observed MW

32kDa

Calculated MW

23kDa

Category

Primary antibody

Applications

WB,IF-P,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC3301

Background

This gene encodes a bifunctional signal transduction molecule. Dopaminergic and glutamatergic receptor stimulation regulates its phosphorylation and function as a kinase or phosphatase inhibitor. As a target for dopamine, this gene may serve as a therapeutic target for neurologic and psychiatric disorders. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:1000 - 1:6000

IF-P 1:200 - 1:1000

IHC-P 1:200 - 1:800

ELISA Recommended starting

concentration is 1 µg/mL.

Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene IDSwiss Prot
84152
Q9UD71

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

DARPP32; DARPP-32

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
$\overline{\Box}$	Т	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: 10 mM sodium HEPES and 150 mM NaCl with proclin300 or sodium azide (as specified on the Certificate of Analysis),100 μ g/ml BSA,50% Glycerol,pH 7.5



Western blot analysis of lysates from Rat brain using DARPP32 Rabbit mAb (A26749) at 1:1000 dilution incubated overnight at 4° C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.








Confocal imaging of paraffin-embedded Mouse brain tissue using DARPP32 Rabbit mAb (A26749, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IF staining. Objective: 40x.




Immunohistochemistry analysis of paraffinembedded Rat brain tissue using DARPP32 Rabbit mAb (A26749) at a dilution of 1\[
\begin{align*}
200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.

Confocal imaging of paraffin-embedded Rat brain tissue using DARPP32 Rabbit mAb (A26749, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IF staining. Objective: 40x.




Immunohistochemistry analysis of paraffinembedded Mouse brain tissue using DARPP32 Rabbit mAb (A26749) at a dilution of 1[200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.

Confocal imaging of paraffin-embedded Rat eyes tissue using DARPP32 Rabbit mAb (A26749, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IF staining. Objective: 40x.




Immunohistochemistry analysis of paraffinembedded Human colon tissue using DARPP32 Rabbit mAb (A26749) at a dilution of 1200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.


Validation Data




Immunohistochemistry analysis of paraffinembedded Mouse colon tissue using DARPP32 Rabbit mAb (A26749) at a dilution of 1 \square 200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse kidney tissue using DARPP32 Rabbit mAb (A26749) at a dilution of 1 \square 200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Rat colon tissue using DARPP32 Rabbit mAb (A26749) at a dilution of $1\square 200$ (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human brain tissue using DARPP32 Rabbit mAb (A26749) at a dilution of 1 \square 200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.