

ABflo® 488 Rabbit anti-Mouse CD50/ICAM-3 mAb

Catalog No.: A26794

Basic Information

Observed MW

Calculated MW

97kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC69845-ABflo488

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Background

Predicted to enable integrin binding activity. Acts upstream of or within phagocytosis. Located in plasma membrane. Is expressed in brain. Orthologous to human ICAM5 (intercellular adhesion molecule 5).

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 15898

Swiss Prot

Q60625

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 32-227 of mouse CD50/ICAM-3 (NP_032345.2).

Synonyms

TLN; CD50; Tlcn; Icam3; ICAM-3; ICAM-5

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	Т	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data





Flow cytometry: 1X10^6 CHO cells (negative control,left) and CHO (Transfection,right) cells were surface-stained with ABflo® 488 Rabbit anti-Mouse CD50/ICAM-3 mAb (A26794,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 CHO (Transfection) cells were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,left) or ABflo® 488 Rabbit anti-Mouse CD50/ICAM-3 mAb (A26794,5 µl/Test,right).