

ABflo® 488 Rabbit anti-Mouse CD50/ICAM-3 mAb

Catalog No.: A26794

Basic Information

Observed MW

Calculated MW

97kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC69845

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Background

Predicted to enable integrin binding activity. Acts upstream of or within phagocytosis. Located in plasma membrane. Is expressed in brain. Orthologous to human ICAM5 (intercellular adhesion molecule 5).

Immunogen Information

Gene ID

15898

Swiss Prot

Q60625

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

TLN; CD50; Tlcn; Icam3; ICAM-3; ICAM-5

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 CHO cells (negative control, left) and CHO (Transfection, right) cells were surface-stained with ABflo® 488 Rabbit anti-Mouse CD50/ICAM-3 mAb (A26794, 5 μ l/Test, orange line) or ABflo® 488 Rabbit IgG isotype control (A22069, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 CHO (Transfection) cells were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069, 5 μ l/Test, left) or ABflo® 488 Rabbit anti-Mouse CD50/ICAM-3 mAb (A26794, 5 μ l/Test, right).