ABflo® 647 Rabbit anti-Mouse CD50/ICAM-3 mAb

Background

(intercellular adhesion molecule 5).

Catalog No.: A26795



Basic Information

Observed MW

Calculated MW 97kDa

Category Primary antibody

Applications FC

Cross-Reactivity Mouse

CloneNo number ARC69845-ABflo647

Conjugate ABflo® 647. Ex:648nm. Em:664nm.

Recommended Dilutions

5 µl per 10^6 cells in

100 µl volume

FC

Immunogen Information

Gene ID 15898 Swiss Prot Q60625

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 32-227 of mouse CD50/ICAM-3 (NP_032345.2).

Predicted to enable integrin binding activity. Acts upstream of or within phagocytosis. Located in plasma membrane. Is expressed in brain. Orthologous to human ICAM5

Synonyms

TLN; CD50; Tlcn; Icam3; ICAM-3; ICAM-5

Contact

6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn


Product Information


Source Rabbit **Isotype** IgG **Purification** Affinity purification

Storage

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.

Validation Data





Flow cytometry: 1X10^6 CHO cells (negative control,left) and CHO (Transfection,right) cells were surface-stained with ABflo® 647 Rabbit anti-Mouse CD50/ICAM-3 mAb (A26795,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line). Flow cytometry: 1X10⁶ CHO (Transfection) cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti-Mouse CD50/ICAM-3 mAb (A26795,5 µl/Test,right).