

ABflo® 700 Rabbit anti-Human CD116 mAb

Catalog No.: A26807

Basic Information

Observed MW

Calculated MW

46kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC59275-ABflo700

Conjugate

ABflo® 700. Ex:690nm. Em:713nm.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Background

The protein encoded by this gene is the alpha subunit of the heterodimeric receptor for colony stimulating factor 2, a cytokine which controls the production, differentiation, and function of granulocytes and macrophages. The encoded protein is a member of the cytokine family of receptors. This gene is found in the pseudoautosomal region (PAR) of the X and Y chromosomes. Multiple transcript variants encoding different isoforms have been found for this gene, with some of the isoforms being membrane-bound and others being soluble.

Immunogen Information

Gene ID

1438

Swiss Prot

P15509

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 23-320 of human CD116 (NP_006131.2).

Synonyms

GMR; CD116; CSF2R; SMDP4; CDw116; CSF2RX; CSF2RY; GMCSFR; CSF2RAX; CSF2RAY; alphaGMR; GMR-alpha; GMCSFR-alpha; GM-CSF-R-alpha

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.


Validation Data



Flow cytometry: 1×10^6 293F cells (negative control, left) and Bewo cells (right) were surface-stained with ABflo® 700 Rabbit anti-Human CD116 mAb (A26807,5 $\mu\text{l}/\text{Test}$, orange line) or ABflo® 700 Rabbit IgG isotype control (A25976,5 $\mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 Human PBMC cells were surface-stained with ABflo® 700 Rabbit anti-Human CD116 mAb (A26807,5 $\mu\text{l}/\text{Test}$, orange line) or ABflo® 700 Rabbit IgG isotype control (A25976,5 $\mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line). Cells in the Monocytes gate were used for analysis.



Flow cytometry: 1×10^6 Bewo cells were surface-stained with ABflo® 700 Rabbit IgG isotype control (A25976,5 $\mu\text{l}/\text{Test}$, left) or ABflo® 700 Rabbit anti-Human CD116 mAb (A26807,5 $\mu\text{l}/\text{Test}$, right).