

ABflo® 610 Rabbit anti-Human β 2 Microglobulin mAb

Catalog No.: A26844

Basic Information

Observed MW

Calculated MW

14kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC60950-ABflo610

Conjugate

ABflo® 610. Ex:421nm. Em:612nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

This gene encodes a serum protein found in association with the major histocompatibility complex (MHC) class I heavy chain on the surface of nearly all nucleated cells. The protein has a predominantly beta-pleated sheet structure that can form amyloid fibrils in some pathological conditions. The encoded antimicrobial protein displays antibacterial activity in amniotic fluid. A mutation in this gene has been shown to result in hypercatabolic hypoproteinemia.

Immunogen Information

Gene ID

567

Swiss Prot

P61769

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 21-119 of human β 2 Microglobulin (NP_004039.1).

Synonyms

IMD43

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.


Validation Data




Flow cytometry: 1×10^6 Daudi cells (negative control, left) and HeLa cells (right) were surface-stained with ABflo® 610 Rabbit anti-Human $\beta 2$ Microglobulin mAb (A26844,5 $\mu\text{l}/\text{Test}$, orange line) or ABflo® 610 Rabbit IgG isotype control (A25826,5 $\mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 HeLa cells were surface-stained with ABflo® 610 Rabbit IgG isotype control (A25826,5 $\mu\text{l}/\text{Test}$, left) or ABflo® 610 Rabbit anti-Human $\beta 2$ Microglobulin mAb (A26844,5 $\mu\text{l}/\text{Test}$, right).



Flow cytometry: 1×10^6 Human PBMC were surface-stained with ABflo® 610 Rabbit anti-Human $\beta 2$ Microglobulin mAb (A26844,5 $\mu\text{l}/\text{Test}$, orange line) or ABflo® 610 Rabbit IgG isotype control (A25826,5 $\mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line). Cells in the Lymphocytes gate were used for analysis.



Flow cytometry: 1×10^6 Human PBMC were surface-stained with ABflo® 610 Rabbit IgG isotype control (A25826,5 $\mu\text{l}/\text{Test}$, left) or ABflo® 610 Rabbit anti-Human $\beta 2$ Microglobulin mAb (A26844,5 $\mu\text{l}/\text{Test}$, right). Cells in the Lymphocytes gate were used for analysis.