

PE Rabbit anti-Mouse CD200/OX2 mAb

Catalog No.: A26859

Basic Information

Observed MW

Calculated MW

31kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC70039-PE

Conjugate

PE. Ex:565nm. Em:574nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

Predicted to enable protein binding activity involved in heterotypic cell-cell adhesion. Involved in negative regulation of cell population proliferation; negative regulation of macrophage activation; and negative regulation of neuroinflammatory response. Located in cell surface. Is expressed in several structures, including anterior naris; aorta; bone; nervous system; and retina. Used to study autoimmune disease. Orthologous to human CD200 (CD200 molecule).

Immunogen Information

Gene ID

17470

Swiss Prot

O54901

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 31-232 of mouse CD200/OX2 (NP_034948.3).

Synonyms

OX2; Mox2

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.


Validation Data




Flow cytometry: 1×10^6 293T cells (negative control, left) and 293T (Transfection, right) cells were surface-stained with PE Rabbit anti-Mouse CD200/OX2 mAb (A26859, 5 μ l/Test, orange line) or PE Rabbit IgG isotype control (A24172, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 RAW 264.7 cells (negative control, left) and BALB/c splenocytes (right) were surface-stained with PE Rabbit anti-Mouse CD200/OX2 mAb (A26859, 5 μ l/Test, orange line) or PE Rabbit IgG isotype control (A24172, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 293T (Transfection) cells were surface-stained with PE Rabbit IgG isotype control (A24172, 5 μ l/Test, left) or PE Rabbit anti-Mouse CD200/OX2 mAb (A26859, 5 μ l/Test, right).



Flow cytometry: 1×10^6 BALB/c splenocytes were surface-stained with PE Rabbit IgG isotype control (A24172, 5 μ l/Test, left) or PE Rabbit anti-Mouse CD200/OX2 mAb (A26859, 5 μ l/Test, right).