

APC Rabbit anti-Human CD38 mAb

Catalog No.: A26866

Basic Information

Observed MW

Calculated MW

13kDa/34kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC5131-01-APC

Conjugate

APC. Ex:650nm. Em:660nm.

Background

The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Immunogen Information

Gene ID

952

Swiss Prot

P28907

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 43-300 of human CD38 (P28907).

Synonyms

ADPRC1; cADPR1; ADPRC 1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.


Validation Data




Flow cytometry: 1×10^6 HepG2 cells (negative control, left) and Daudi cells (right) were surface-stained with APC Rabbit anti-Human CD38 mAb (A26866, $5 \mu\text{l}/\text{Test}$, orange line) or APC Rabbit IgG isotype control (A24173, $5 \mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 Daudi cells were surface-stained with APC Rabbit IgG isotype control (A24173, $5 \mu\text{l}/\text{Test}$, left) or APC Rabbit anti-Human CD38 mAb (A26866, $5 \mu\text{l}/\text{Test}$, right).



Flow cytometry: 1×10^6 Human PBMC were surface-stained with APC Rabbit anti-Human CD38 mAb (A26866, $5 \mu\text{l}/\text{Test}$, orange line) or APC Rabbit IgG isotype control (A24173, $5 \mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 Human PBMC cells were surface-stained with APC Rabbit IgG isotype control (A24173, $5 \mu\text{l}/\text{Test}$, left) or APC Rabbit anti-Human CD38 mAb (A26866, $5 \mu\text{l}/\text{Test}$, right).