ABclonal www.abclonal.com

ABflo® 488 Rabbit anti-Human P4HB mAb

Catalog No.: A26906

Basic Information

Observed MW

Calculated MW

57kDa

Category

Primary antibody

Applications

FC (intra)

Cross-Reactivity

Human

CloneNo number

ARC70670

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Background

This gene encodes the beta subunit of prolyl 4-hydroxylase, a highly abundant multifunctional enzyme that belongs to the protein disulfide isomerase family. When present as a tetramer consisting of two alpha and two beta subunits, this enzyme is involved in hydroxylation of prolyl residues in preprocollagen. This enzyme is also a disulfide isomerase containing two thioredoxin domains that catalyze the formation, breakage and rearrangement of disulfide bonds. Other known functions include its ability to act as a chaperone that inhibits aggregation of misfolded proteins in a concentration-dependent manner, its ability to bind thyroid hormone, its role in both the influx and efflux of S-nitrosothiol-bound nitric oxide, and its function as a subunit of the microsomal triglyceride transfer protein complex.

Recommended Dilutions

FC (intra)

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 5034 **Swiss Prot**

P07237

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

DSI; GIT; PDI; PHDB; PDIA1; PO4DB; PO4HB; PROHB; CLCRP1; ERBA2L; P4Hbeta

Contact

a	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data









Flow cytometry: 1X10^6 knockout (KO) HeLa cells (negative control,left) and HeLa cells (right) were intracellularly-stained with ABflo® 488 Rabbit anti-Human P4HB mAb (A26906,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 HeLa cells were intracellularly-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,left) or ABflo® 488 Rabbit anti-Human P4HB mAb (A26906,5 µl/Test,right).