

APC/Cyanine7 Rabbit anti-Human/Monkey CD8a mAb

Catalog No.: A26941

Basic Information

Observed MW

Calculated MW

21kDa/25kDa/30kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human, Cynomolgus

CloneNo number

ARC55248

Conjugate

APC-Cy7. Ex:651nm. Em:779nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen acts as a coreceptor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell in the context of class I MHC molecules. The coreceptor functions as either a homodimer composed of two alpha chains or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 alpha chain. Multiple transcript variants encoding different isoforms have been found for this gene. The major protein isoforms of this gene differ by the presence or absence of a transmembrane domain and thus differ in being a membrane-anchored or secreted protein.

Immunogen Information

Gene ID

925

Swiss Prot

P01732

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD8; p32; Leu2; CD8alpha

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 Cynomolgus PBMC were surface-stained with ABflo® 488 Rabbit anti-Human/Monkey CD3 mAb (A26283, 5 μ l/Test) and APC/Cyanine7 Rabbit IgG isotype control (5 μ l/Test, left) or APC/Cyanine7 Rabbit anti-Human/Monkey CD8a mAb (A26941, 5 μ l/Test, right). Cells in the lymphocyte gate were used for analysis.

Flow cytometry: 1×10^6 Human PBMC were surface-stained with ABflo® 488 Rabbit anti-Human/Monkey CD3 mAb (A26283, 5 μ l/Test) and APC/Cyanine7 Rabbit IgG isotype control (5 μ l/Test, left) or APC/Cyanine7 Rabbit anti-Human/Monkey CD8a mAb (A26941, 5 μ l/Test, right). Cells in the lymphocyte gate were used for analysis.