

ANPEP/CD13 Rabbit mAb

Catalog No.: A26949 **Recombinant**

Basic Information

Observed MW

160kDa

Calculated MW

109kDa

Category

Primary antibody

Applications

WB, IF-P, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC69826

Background

Predicted to enable metalloaminopeptidase activity; peptide binding activity; and zinc ion binding activity. Predicted to be involved in several processes, including negative regulation of renal sodium excretion; peptide catabolic process; and proteolysis. Predicted to act upstream of or within angiogenesis and cell differentiation. Located in external side of plasma membrane. Is expressed in several structures, including alimentary system; brain; metanephros; reproductive system; and sensory organ. Orthologous to human ANPEP (alanyl aminopeptidase, membrane).

Recommended Dilutions

WB 1:17000 - 1:100000**IF-P** 1:200 - 1:800**IHC-P** 1:1000 - 1:10000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

16790

Swiss Prot

P97449

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Apn; AP-M; AP-N; Cd13; P150; CD13

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage




Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data






Western blot analysis of various lysates using ANPEP/CD13 Rabbit mAb (A26949) at 1:17000 dilution incubated overnight at 4°C.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Negative control (NC): 293T
 Exposure time: 30s.



Confocal imaging of paraffin-embedded Mouse kidney tissue using ANPEP/CD13 Rabbit mAb (A26949, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Confocal imaging of paraffin-embedded Mouse spleen tissue using ANPEP/CD13 Rabbit mAb (A26949, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Confocal imaging of paraffin-embedded Rat kidney tissue using ANPEP/CD13 Rabbit mAb (A26949, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Rat spleen tissue using ANPEP/CD13 Rabbit mAb (A26949, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.


Confocal imaging of paraffin-embedded Human kidney tissue using ANPEP/CD13 Rabbit mAb (A26949, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using ANPEP/CD13 Rabbit mAb (A26949) at a dilution of 1:9000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.


Validation Data




Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using ANPEP/CD13 Rabbit mAb (A26949) at a dilution of 1:9000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using ANPEP/CD13 Rabbit mAb (A26949) at a dilution of 1:9000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat kidney tissue using ANPEP/CD13 Rabbit mAb (A26949) at a dilution of 1:9000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using ANPEP/CD13 Rabbit mAb (A26949) at a dilution of 1:9000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.