

GDNF Receptor alpha 1/GFRA1 Rabbit mAb

Catalog No.: A27002 **Recombinant** **1 Publications**

Basic Information

Observed MW

Refer to figures

Calculated MW

51kDa

Category

Primary antibody

Applications

IF/ICC, IF-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC59643

Background

This gene encodes a member of the glial cell line-derived neurotrophic factor receptor (GDNFR) family of proteins. The encoded preproprotein is proteolytically processed to generate the mature receptor. Glial cell line-derived neurotrophic factor (GDNF) and neurturin (NTN) are two structurally related, potent neurotrophic factors that play key roles in the control of neuron survival and differentiation. This receptor is a glycosylphosphatidylinositol (GPI)-linked cell surface receptor for both GDNF and NTN, and mediates activation of the RET tyrosine kinase receptor. This gene is a candidate gene for Hirschsprung disease. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed.

Recommended Dilutions

IF/ICC 1:100 - 1:400**IF-P** 1:100 - 1:400

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

2674

Swiss Prot

P56159

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

GDNFR; RET1L; RETL1; RHDA4; TRNR1; GDNFRA; GFRalpha-1; GFR-ALPHA-1; GDNFR-alpha-1

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG


Purification

Affinity purification


Storage

Store at -20°C


Validation Data



Immunofluorescence analysis of MCF7 cells using GDNF Receptor alpha 1/GFRA1 Rabbit mAb (A27002) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Mouse brain tissue using GDNF Receptor alpha 1/GFRA1 Rabbit mAb (A27002) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining. High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining.



Immunofluorescence analysis of Rat brain tissue using GDNF Receptor alpha 1/GFRA1 Rabbit mAb (A27002) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining. High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining.