ABclonal www.abclonal.com

ABflo® 647 Rabbit anti-Mouse CD270/HVEM mAb

Catalog No.: A27136

Basic Information

Observed MW

Calculated MW

30kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC71587-ABflo647

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Background

Predicted to enable cytokine binding activity and ubiquitin protein ligase binding activity. Acts upstream of or within several processes, including defense response to bacterium; negative regulation of alpha-beta T cell proliferation; and positive regulation of macromolecule metabolic process. Located in external side of plasma membrane. Is expressed in liver lobe. Orthologous to human TNFRSF14 (TNF receptor superfamily member 14).

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 230979

Swiss Prot Q80WM9

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 39-207 of mouse CD270/HVEM (NP_849262.1).

Synonyms

TR2; Atar; HveA; Hvem; Tnfrs14

Contact

2		400-999-6126
\bowtie	Τ	cn.market@abclonal.com.cn
•	Т	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data









Flow cytometry: 1X10^6 C2C12 cells (negative control,left) and C57BL/6 mouse splenocytes (right) were surface-stained with ABflo® 647 Rabbit anti-Mouse CD270/HVEM mAb (A27136,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Nonfluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 C57BL/6 mouse splenocytes were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti-Mouse CD270/HVEM mAb (A27136,5 µl/Test,right).