# APC Rat anti-Mouse CD86 mAb

Catalog No.: A27138



### **Basic Information**

#### **Observed MW**

Calculated MW

35kDa

Category

Primary antibody

**Applications** 

FC

**Cross-Reactivity** 

Mouse

CloneNo number

ATC0032-APC

Conjugate

APC. Ex:650nm. Em:660nm.

# **Background**

Predicted to enable signaling receptor binding activity. Involved in several processes, including CD40 signaling pathway; activation of phospholipase C activity; and positive regulation of macromolecule metabolic process. Acts upstream of or within several processes, including cellular response to lipopolysaccharide; defense response to virus; and positive regulation of T cell proliferation. Located in external side of plasma membrane and intracellular membrane-bounded organelle. Is expressed in central nervous system and retina. Used to study Guillain-Barre syndrome. Human ortholog(s) of this gene implicated in several diseases, including Henoch-Schoenlein purpura; autoimmune disease (multiple); chronic lymphocytic leukemia; chronic obstructive pulmonary disease; and systemic scleroderma. Orthologous to human CD86 (CD86 molecule).

## **Recommended Dilutions**

FC

5  $\mu$ l per 10^6 cells in 100  $\mu$ l volume

### **Immunogen Information**

**Gene ID** 12524

**Swiss Prot** 

P42082

**Immunogen** 

Recombinant protein of mouse CD86.

**Synonyms** 

B7; B70; MB7; B7-2; B7.2; CLS1; Ly58; ETC-1; Ly-58; MB7-2; Cd28l2; TS/A-2; CD86

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	Т	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRatIgGAffinity purification

**Storage** 

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

# **Validation Data**



Flow cytometry:  $1X10^6$  C57BL/6 mouse splenocytes were surface-stained with APC Rat IgG isotype control (2  $\mu$ g/Test,left) or APC Rat anti-Mouse CD86 mAb (A27138,5  $\mu$ l/Test,right).