

E2F6 Rabbit pAb

Catalog No.: A2718

2 Publications

Basic Information

Observed MW

35kDa

Calculated MW

32kDa

Category

Primary antibody

Applications

WB, IF/ICC, IP, ELISA, ChIP

Cross-Reactivity

Human, Mouse

Background

This gene encodes a member of a family of transcription factors that play a crucial role in the control of the cell cycle. The protein encoded by this gene lacks the transactivation and tumor suppressor protein association domains found in other family members, and contains a modular suppression domain that functions in the inhibition of transcription. It interacts in a complex with chromatin modifying factors. There are pseudogenes for this gene on chromosomes 22 and X. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:500 - 1:2000**IF/ICC** 1:50 - 1:200**IP** 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.**ChIP** 5µg antibody for
10µg-15µg of Chromatin

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn

Immunogen Information

Gene ID

1876

Swiss Prot

O75461

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-281 of human E2F6 (NP_937987.2).

Synonyms

E2F-6; E2F6

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.




www.abclonal.com.cn


Validation Data



Western blot analysis of lysates from Mouse lung using E2F6 Rabbit pAb (A2718) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.




Immunoprecipitation analysis of 200 µg extracts of MCF-7 cells using 3 µg E2F6 antibody (A2718). Western blot was performed from the immunoprecipitate using E2F6 antibody (A2718) at a dilution of 1:500.



Chromatin immunoprecipitation analysis of extracts of 293T cells, using E2F6 antibody (A2718) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.

Validation Data



Immunofluorescence analysis of U2OS cells using E2F6 Rabbit pAb (A2718). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.