PE Rabbit anti-Human CD18 mAb

PE Nappit aliti-Hulliali CD10 IIIAL



Catalog No.: A27214

Basic Information

Observed MW

Calculated MW

85kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC72097-PE

Conjugate

PE. Ex:565nm. Em:574nm.

Background

This gene encodes an integrin beta chain, which combines with multiple different alpha chains to form different integrin heterodimers. Integrins are integral cell-surface proteins that participate in cell adhesion as well as cell-surface mediated signalling. The encoded protein plays an important role in immune response and defects in this gene cause leukocyte adhesion deficiency. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 3689 **Swiss Prot**

P05107

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-769 of human CD18 (NP_000202.3).

Synonyms

LAD; CD18; MF17; MFI7; LCAMB; LFA-1; MAC-1

Contact

2		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data





Flow cytometry: 1X10^6 HeLa cells (negative control,left) and THP-1 cells (right) were surface-stained with PE Rabbit anti-Human CD18 mAb (A27214,5 μ l/Test,orange line) or PE Rabbit IgG isotype control (A24172,5 μ l/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: $1X10^6$ Human PBMC were surface-stained with PE Rabbit IgG isotype control (A24172,5 μ I/Test,Ieft) or PE Rabbit anti-Human CD18 mAb (A27214,5 μ I/Test,right).