

Acetyl-Histone H3-K14 Rabbit mAb

Catalog No.: A27265 **Recombinant**

Basic Information

Observed MW

17kDa

Calculated MW

16kDa

Category

Primary antibody

Applications

WB,DB,CHIP,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC70064

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB 1:11000 - 1:66000**DB** 1:1000 - 1:2000**CHIP** 3µg antibody for
10µg-15µg of chromatin**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431


Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data




Western blot analysis of lysates from C6 cells using Acetyl-Histone H3-K14 Rabbit mAb (A27265) at 1:11000 dilution incubated overnight at 4°C. C6 cells were treated with TSA (1 μ M) at 37°C for 18 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.




Western blot analysis of lysates from HeLa cells using Acetyl-Histone H3-K14 Rabbit mAb (A27265) at 1:11000 dilution incubated overnight at 4°C. HeLa cells were treated with TSA (1 μ M) at 37°C for 18 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.



Western blot analysis of lysates from NIH/3T3 cells using Acetyl-Histone H3-K14 Rabbit mAb (A27265) at 1:11000 dilution incubated overnight at 4°C. NIH/3T3 cells were treated with Sodium lactate (100 mM) for 24h.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 30 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.

Validation Data



Chromatin immunoprecipitation was performed with 10 µg of cross-linked chromatin from HeLa, using 3 µg of Acetyl-Histone H3-K14 Rabbit mAb (A27265) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.



Dot-blot analysis of all sorts of peptides using Acetyl-Histone H3-K14 Rabbit mAb (A27265) at 1:1000 dilution.