[KD Validated] MCT1 Rabbit mAb

Catalog No.: A27270 Recombinant



Basic Information

Observed MW

43kDa

Calculated MW

54kDa

Category

Primary antibody

Applications

WB,IF/ICC,IF-P,IHC-P,ELISA

Cross-Reactivity

Human

CloneNo number

ARC72781

Background

The protein encoded by this gene is a proton-linked monocarboxylate transporter that catalyzes the movement of many monocarboxylates, such as lactate and pyruvate, across the plasma membrane. Mutations in this gene are associated with erythrocyte lactate transporter defect. Alternatively spliced transcript variants have been found for this gene.

Recommended Dilutions

WB	1:2000 - 1:8000
IF/ICC	1:200 - 1:800
IF-P	1:200 - 1:800

ELISA Recommended starting

concentration is 1 µg/mL.

Please optimize the
concentration based on
your specific assay
requirements.

1:200 - 1:800

Immunogen Information

Gene ID	Swiss Prot
6566	P53985

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

MCT; HHF7; MCT1; MCT1D

Contact

IHC-P

a	400-999-6126
×	cn.market@abclonal.com.cn
	www.abclonal.com.cn


Product Information

Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.





Western blot analysis of lysates from wild type (WT) and MCT1 knockdown (KD) 293T cells using [KD Validated] MCT1 Rabbit mAb (A27270) at 1:2000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.









Confocal imaging of HeLa cells using [KD Validated] MCT1 Rabbit mAb (A27270, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with $\alpha\text{-Tubulin}$ Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.

Confocal imaging of paraffin-embedded Human colon cancer tissue using [KD Validated] MCT1 Rabbit mAb (A27270, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Immunohistochemistry analysis of paraffinembedded Human colon carcinoma tissue using [KD Validated] MCT1 Rabbit mAb (A27270) at a dilution of 1:600 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human esophagus tissue using [KD Validated] MCT1 Rabbit mAb (A27270) at a dilution of 1:600 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human liver cancer tissue using [KD Validated] MCT1 Rabbit mAb (A27270) at a dilution of 1:600 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.