

# PerCP Rabbit anti-Mouse CD45 mAb

Catalog No.: A27305

## Basic Information

### Observed MW

### Calculated MW

130kDa/140kDa/144kDa

### Category

Primary antibody

### Applications

FC

### Cross-Reactivity

Mouse

### CloneNo number

ARC60962

### Conjugate

PerCP. Ex:482nm. Em:678nm.

## Background

Enables several functions, including heparan sulfate proteoglycan binding activity; heparin binding activity; and protein tyrosine phosphatase activity. Involved in several processes, including lymphocyte differentiation; positive regulation of macromolecule metabolic process; and regulation of signal transduction. Acts upstream of or within several processes, including lymphocyte differentiation; positive regulation of lymphocyte activation; and regulation of protein phosphorylation. Located in external side of plasma membrane; focal adhesion; and membrane raft. Is expressed in several structures, including 3rd branchial arch; alimentary system; cardiovascular system; hemolymphoid system; and placenta. Used to study systemic lupus erythematosus. Human ortholog(s) of this gene implicated in hepatitis C; multiple sclerosis; severe combined immunodeficiency; and severe combined immunodeficiency, autosomal recessive, T cell-negative, B cell-positive, Nk cell-positive. Orthologous to human PTPRC (protein tyrosine phosphatase receptor type C).

## Recommended Dilutions

FC 5 µl per 10<sup>6</sup> cells in  
100 µl volume

## Immunogen Information

### Gene ID

19264

### Swiss Prot

P06800

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

loc; B220; Cd45; L-CA; Ly-5; T200; CD45R; Lyt-4

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

## Validation Data



Flow cytometry:  $1 \times 10^6$  C2C12 cells (negative control, left) and C57BL/6 mouse splenocytes (right) were surface-stained with PerCP Rabbit anti-Mouse CD45 mAb (A27305, 5  $\mu$ l/Test, orange line) or PerCP Rabbit IgG isotype control (A24204, 5  $\mu$ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:  $1 \times 10^6$  C57BL/6 mouse splenocytes were surface-stained with PerCP Rabbit IgG isotype control (A24204, 5  $\mu$ l/Test, left) or PerCP Rabbit anti-Mouse CD45 mAb (A27305, 5  $\mu$ l/Test, right).