PE Rabbit anti-Human Calreticulin mAb

ABclonal www.abclonal.com

Catalog No.: A27386

Basic Information

Observed MW

Calculated MW

48kDa

Category

Primary antibody

Applications

FC (intra)

Cross-Reactivity

Human

CloneNo number

ARC71596

Conjugate

PE. Ex:565nm. Em:574nm.

Background

Calreticulin is a highly conserved chaperone protein which resides primarily in the endoplasmic reticulum, and is involved in a variety of cellular processes, among them, cell adhesion. Additionally, it functions in protein folding quality control and calcium homeostasis. Calreticulin is also found in the nucleus, suggesting that it may have a role in transcription regulation. Systemic lupus erythematosus is associated with increased autoantibody titers against calreticulin. Recurrent mutations in calreticulin have been linked to various neoplasms, including the myeloproliferative type.

Recommended Dilutions

FC (intra)

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 811 **Swiss Prot**

P27797

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

RO; CRT; SSA; cC1qR; HEL-S-99n

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	T	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data









Flow cytometry: 1×10^6 knockout (KO) HeLa cells (negative control,left) and HeLa cells (right) were intracellularly-stained with PE Rabbit anti-Human Calreticulin mAb (A27386,5 μ I/Test,orange line) or PE Rabbit IgG isotype control (A24172,5 μ I/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: $1X10^6$ HeLa cells were intracellularly-stained with PE Rabbit IgG isotype control (A24172,5 μ I/Test,left) or PE Rabbit anti-Human Calreticulin mAb (A27386,5 μ I/Test,right).