

PE-CF594 Rabbit anti-Human/Monkey HLA-DR mAb

Catalog No.: A27408

Basic Information

Observed MW

Calculated MW
29kDa

Category
Primary antibody

Applications
FC

Cross-Reactivity
Human, Cynomolgus

CloneNo number
ARC5141-01

Conjugate
PE-CF594. Ex:566nm. Em:615nm.

Background

HLA-DRA is one of the HLA class II alpha chain paralogues. This class II molecule is a heterodimer consisting of an alpha and a beta chain, both anchored in the membrane. This molecule is expressed on the surface of various antigen presenting cells such as B lymphocytes, dendritic cells, and monocytes/macrophages, and plays a central role in the immune system and response by presenting peptides derived from extracellular proteins, in particular, pathogen-derived peptides to T cells. The alpha chain is approximately 33-35 kDa and its gene contains 5 exons. Exon 1 encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, and exon 4 encodes the transmembrane domain and the cytoplasmic tail. DRA does not have polymorphisms in the peptide binding part and acts as the sole alpha chain for DRB1, DRB3, DRB4 and DRB5.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Immunogen Information

Gene ID
3122

Swiss Prot
P01903

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms
HLA-DRA1

Contact

☎ | 400-999-6126
✉ | cn.market@abclonal.com.cn
🌐 | www.abclonal.com.cn

Product Information

Source
Rabbit


Isotype
IgG

Purification
Affinity purification


Storage

Store at 2-8°C. Avoid freeze.
Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.



Validation Data






Flow cytometry: 1×10^6 PC-3 cells (negative control, left) and Human PBMC (right) were surface-stained with PE-CF594 Rabbit anti-Human HLA-DR mAb (A27408, 5 μ l/Test, orange line) or PE-CF594 Rabbit IgG isotype control (5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line). Cells in the Lymphocytes gate were used for analysis.



Flow cytometry: 1×10^6 Cynomolgus PBMC were surface-stained with PE-CF594 Rabbit IgG isotype control (5 μ l/Test, left) or PE-CF594 Rabbit anti-Human/Monkey HLA-DR mAb (A27408, 5 μ l/Test, right). Cells in the lymphocyte gate were used for analysis.



Flow cytometry: 1×10^6 Human PBMC were surface-stained with PE-CF594 Rabbit IgG isotype control (5 μ l/Test, left) or PE-CF594 Rabbit anti-Human/Monkey HLA-DR mAb (A27408, 5 μ l/Test, right). Cells in the lymphocyte gate were used for analysis.



Flow cytometry: 1×10^6 Human PBMC were surface-stained with ABflo® 488 Rabbit anti-Human/Monkey CD3 mAb (A26283, 5 μ l/Test) and PE-CF594 Rabbit IgG isotype control (5 μ l/Test, left) or PE-CF594 Rabbit anti-Human/Monkey HLA-DR mAb (A27408, 5 μ l/Test, right). Cells in the lymphocyte gate were used for analysis.