

# Acetyl-Histone H3-K27 Rabbit mAb

Catalog No.: A2771   **Recombinant**   **6 Publications**

## Basic Information

### Observed MW

17kDa

### Calculated MW

15kDa

### Category

Primary antibody

### Applications

WB, DB, IHC-P, IF/ICC, IP, ELISA, ChIP

### Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range  
Predicted)

### Clone/No. number

ARC53670

## Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

## Recommended Dilutions

**WB**                    1:10000 - 1:120000

**DB**                    1:500 - 1:2000

**IHC-P**                1:500 - 1:1000

**IF/ICC**               1:50 - 1:200

**IP**                    0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells

**ELISA**                Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

**ChIP**                2µg antibody for  
5µg-10µg of Chromatin

## Immunogen Information

### Gene ID

8290/8350

### Swiss Prot

Q16695/P68431

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## **Contact**

---


-  | 400-999-6126
-  | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)
-  | [www.abclonal.com.cn](http://www.abclonal.com.cn)

---


## Validation Data




Immunoprecipitation analysis of 600 µg extracts of HeLa cells treated with TSA using 5 µg Acetyl-Histone H3-K27 Rabbit mAb(A2771). Western blot was performed from the immunoprecipitate using Acetyl-Histone H3-K27 antibody (A2771) at a dilution of 1:50000.




Immunoprecipitation of Acetyl-Histone H3-K27 from 600 µg extracts of NIH/3T3 cells treated with TSA(1µM) 18h was performed using 2 µg of Acetyl-Histone H3-K27 Rabbit mAb (A2771). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X non-reducing Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1 : 50000.




## Validation Data




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Acetyl-Histone H3-K27 antibody (A2771) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.




Dot-blot analysis of all sorts of peptides using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at 1:200000 dilution.




Immunofluorescence analysis of HeLa treated with TSA and HeLa cells using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human liver tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human small intestine tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human thyroid cancer tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.


## Validation Data




Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.