

APC-ABflo® 700 Rabbit anti-Mouse CD44 mAb

Catalog No.: A27729

Basic Information

Observed MW

Calculated MW

86kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC63783

Conjugate

APC-ABflo® 700. Ex:633nm. Em:715nm.

Background

Enables hyaluronic acid binding activity and type II transforming growth factor beta receptor binding activity. Contributes to cytokine binding activity and cytokine receptor activity. Involved in several processes, including negative regulation of T cell activation; positive regulation of protein phosphorylation; and regulation of intracellular signal transduction. Acts upstream of or within several processes, including Wnt signaling pathway; morphogenesis of a branching epithelium; and wound healing involved in inflammatory response. Located in basolateral plasma membrane; external side of plasma membrane; and microvillus. Part of macrophage migration inhibitory factor receptor complex. Is expressed in several structures, including alimentary system; branchial arch; central nervous system; genitourinary system; and limb. Human ortholog(s) of this gene implicated in breast carcinoma (multiple); carcinoma (multiple); and prostate cancer. Orthologous to human CD44 (CD44 molecule (Indian blood group)).

Recommended Dilutions

FC ≤ 0.25 μ g per million cells
in 100 μ l volume

Immunogen Information

Gene ID

12505

Swiss Prot

P15379

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 23-176 of mouse CD44 (NP_033981.2).

Synonyms

Ly-24; Pgp-1; HERMES

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG


Purification

Affinity purification

Storage

Store at 2-8°C. Avoid freeze.
Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.

Validation Data



Flow cytometry: 1X10⁶ C57BL/6 mouse splenocytes were surface-stained with PE Rabbit anti-Mouse CD62L mAb (A26715, 5 μ l/Test) and APC-ABflo® 700 Rabbit IgG isotype control (0.25 μ g, left) or APC-ABflo® 700 Rabbit anti-Mouse CD44 mAb (A27729, 0.25 μ g, right). Cells in the CD4 positive gate were used for analysis.