

APC/Cyanine7 Rabbit anti-Human/Monkey CD38 PolymAb®

Catalog No.: A28063PM

Basic Information

Observed MW

Calculated MW

34kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human, Cynomolgus

Conjugate

APC-Cy7. Ex:651nm. Em:779nm.

Background

The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 952 **Swiss Prot**

P28907

Immunogen

This information is considered to be commercially sensitive.

Synonyms

ADPRC1; cADPR1; ADPRC 1

Contact

2		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1X10^6 Human PBMC were surface-stained with APC/Cyanine7 Rabbit IgG isotype control (5 µl/Test,left) or APC/Cyanine7 Rabbit anti-Human CD38 polymAb® (A28063PM,5 µl/Test,right). Cells in the lymphocyte gate were used for analysis.