ABclonal www.abclonal.com

ABflo® 450 Rabbit anti-Human CD68 mAb

Catalog No.: A28159

Basic Information

Observed MW

Calculated MW

37 kDa

Category

Primary antibody

Applications

FC (intra)

Cross-Reactivity

Human

CloneNo number

ARC56633

Conjugate

ABflo® 450. Ex:406nm. Em:445nm.

Background

This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. It is a type I integral membrane protein with a heavily glycosylated extracellular domain and binds to tissue- and organ-specific lectins or selectins. The protein is also a member of the scavenger receptor family. Scavenger receptors typically function to clear cellular debris, promote phagocytosis, and mediate the recruitment and activation of macrophages. Alternative splicing results in multiple transcripts encoding different isoforms.

Recommended Dilutions

FC (intra)

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 968 **Swiss Prot**

P34810

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

GP110; LAMP4; SCARD1

Contact

2		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry:1X10^6 Human PBMC were surface-stained with PE Rabbit anti-Human/Monkey CD14 mAb (A25938,5 µl/Test) and then intracellularly-stained with ABflo® 450 Rabbit IgG isotype control (A27451,5 µl/Test,left) or ABflo® 450 Rabbit anti-Human CD68 mAb (A28159,5 µl/Test,right). Cells in the monocyte gate were used for analysis.