

# Biotin Rat anti-Mouse CD45R/B220 mAb

Catalog No.: A28202

## Basic Information

### Observed MW

### Calculated MW

130kDa/140kDa/144kDa

### Category

Primary antibody

### Applications

FC

### Cross-Reactivity

Mouse

### CloneNo number

ATC10005

### Conjugate

Biotin

## Recommended Dilutions

**FC** ≤0.25 µg per million cells  
in 100 µl volume

## Background

Enables several functions, including heparan sulfate proteoglycan binding activity; heparin binding activity; and protein tyrosine phosphatase activity. Involved in several processes, including lymphocyte differentiation; positive regulation of macromolecule metabolic process; and regulation of signal transduction. Acts upstream of or within several processes, including lymphocyte differentiation; positive regulation of lymphocyte activation; and regulation of protein phosphorylation. Located in external side of plasma membrane; focal adhesion; and membrane raft. Is expressed in several structures, including 3rd branchial arch; alimentary system; cardiovascular system; hemolymphoid system; and placenta. Used to study systemic lupus erythematosus. Human ortholog(s) of this gene implicated in hepatitis C; multiple sclerosis; severe combined immunodeficiency; and severe combined immunodeficiency, autosomal recessive, T cell-negative, B cell-positive, Nk cell-positive. Orthologous to human PTPRC (protein tyrosine phosphatase receptor type C).

## Immunogen Information

### Gene ID

19264/5788

### Swiss Prot

P06800/P08575

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

loc; B220; Cd45; L-CA; Ly-5; T200; CD45R; Lyt-4

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rat

### Isotype

Rat / IgG2a, kappa

### Purification

Affinity purification


### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

## Validation Data

---



Flow cytometry: 1X10<sup>6</sup> C57BL/6 mouse splenocytes were surface-stained with Biotin Rat IgG2a isotype control (5 µl/Test,left) or Biotin Rat anti-Mouse CD45R/B220 mAb (A28202,0.25 µg,right), followed by ABflo® 488 Streptavidin staining.